

A Note on Admissible Mannheim Curves in Galilean Space G_3

S. ERSOY, M. AKYİĞİT, M. TOSUN

sersoy@sakarya.edu.tr, makyigit@sakarya.edu.tr, tosun@sakarya.edu.tr
Department of Mathematics, Faculty of Arts and Sciences,
Sakarya University, Sakarya, 54187 TURKEY

March 17, 2010

Abstract

The aim of this paper is to study the Mannheim partner curves in three dimensional Galilean space G_3 . Some well known theorems are obtained related to Mannheim curves.

Mathematics Subject Classification (2010). 53A35, 51M30.

Keywords: Galilean space, Mannheim curve, admissible curve.

1 Introduction

In 1850 J. Bertrand defined Bertrand curves, with the help of this definition Liu and Wang called the Mannheim pair which the principal normal vector of the first curve coincides with the binormal vector of the second curve and they obtained the necessary and sufficient conditions between the curvature and the torsions for a curve to be the Mannheim partner curves, [1]. Moreover, Orbay and Kasap examined on Mannheim curves in $\mathbb{E}^3$, [2]. The geometry of the Galilean space G_3 has been treated in detail in Röschel's habilitation, [3]. Furthermore, Kamenarovic and Sipus studied about Galilean space, [4, 5]. The properties of the curves in the Galilean space are studied in [6], [7], [8]. In this paper, we gave some theorems and relations about the curvatures and torsions of admissible Mannheim curves in 3-dimensional Galilean space G_3 .

2 Preliminaries

The Galilean space G_3 is a Cayley-Klein space equipped with the projective metric of signature $(0,0,+,+)$, as in [9]. The absolute figure of the Galilean Geometry consist of an ordered triple $\{w, f, I\}$, where w is the ideal (absolute) plane, f is the line (absolute line) in w and I is the fixed elliptic involution of

points of f , [4]. In the non-homogeneous coordinates the similarity group H_8 has the form

$$\begin{aligned}\bar{x} &= a_{11} + a_{12}x \\ \bar{y} &= a_{21} + a_{22}x + a_{23}y \cos \varphi + a_{23}z \sin \varphi \\ \bar{z} &= a_{31} + a_{32}x - a_{23}y \sin \varphi + a_{23}z \cos \varphi\end{aligned}\quad (2.1)$$

where a_{ij} and φ are real numbers, [6].

In what follows the coefficients a_{12} and a_{23} will play the special role. In particular, for $a_{12} = a_{23} = 1$, (2.1) defines the group $B_6 \subset H_8$ of isometries of Galilean space G_3 .

In G_3 there are four classes of lines:

- i)* (proper) non-isotropic lines- they don't meet the absolute line f .
- ii)* (proper) isotropic lines- lines that don't belong to the plane w but meet the absolute line f .
- iii)* unproper non-isotropic lines-all lines of w but f .
- iv)* the absolute line f .

Planes $x = \text{constant}$ are Euclidean and so is the plane w . Other planes are isotropic, [5].

Galilean scalar product can be written as

$$\langle u_1, u_2 \rangle = \begin{cases} x_1 x_2 & , \text{ if } x_1 \neq 0 \vee x_2 \neq 0 \\ y_1 y_2 + z_1 z_2 & , \text{ if } x_1 = 0 \wedge x_2 = 0 \end{cases}$$

where $u_1 = (x_1, y_1, z_1)$ and $u_2 = (x_2, y_2, z_2)$. It leaves invariant the Galilean norm of the vector $u = (x, y, z)$ defined by

$$\|u\| = \begin{cases} x & , x \neq 0 \\ \sqrt{y^2 + z^2} & , x = 0. \end{cases}$$

Let α be a curve given in the coordinate form

$$\begin{aligned}\alpha : I &\rightarrow G_3, & I &\subset \mathbb{R} \\ t &\rightarrow \alpha(t) = (x(t), y(t), z(t))\end{aligned}\quad (2.2)$$

where $x(t), y(t), z(t) \in C^3$ and t is a real interval. If $x'(t) \neq 0$, then the curve α is called admissible curve.

Let α be an admissible curve in G_3 , is parameterized by arc length s , is given by

$$\alpha(s) = (s, y(s), z(s))$$

where the curvature $\kappa(s)$ and the torsion $\tau(s)$ are

$$\kappa(s) = \sqrt{y''^2(s) + z''^2(s)} \quad \text{and} \quad \tau(s) = \frac{\det[\alpha'(s), \alpha''(s), \alpha'''(s)]}{\kappa^2(s)}, \quad (2.3)$$

respectively. The associated moving Frenet frame is

$$\begin{aligned}T(s) &= \alpha'(s) = (1, y'(s), z'(s)) \\ N(s) &= \frac{1}{\kappa(s)} \alpha''(s) = \frac{1}{\kappa(s)} (0, y''(s), z''(s)) \\ B(s) &= \frac{1}{\kappa(s)} (0, -z''(s), y''(s)).\end{aligned}\quad (2.4)$$

Here T, N and B are called the tangent vector, principal normal vector and binormal vector fields of the curve α , respectively. Then for the curve α , the following Frenet equations are given by

$$\begin{aligned} T'(s) &= \kappa(s)N(s) \\ N'(s) &= \tau(s)B(s) \\ B'(s) &= -\tau(s)N(s) \end{aligned} \quad (2.5)$$

where T, N, B are mutually orthogonal vectors, [5].

3 Admissible Mannheim Curves in Galilean Space G_3

In this section, we defined the admissible Mannheim curve and gave some theorems related to these curves in G_3 .

Definition 3.1 Let α and α^* be an admissible curves with the Frenet frames along $\{T, N, B\}$ and $\{T^*, N^*, B^*\}$, respectively. The curvature and torsion of α and α^* , respectively, $\kappa(s), \tau(s)$ and $\kappa^*(s), \tau^*(s)$ never vanish for all $s \in I$ in G_3 . If the principal normal vector field N of α coincidence with the binormal vector field B^* of α^* at the corresponding points of the admissible curves α and α^* . Then α is called an admissible Mannheim curve and α^* is an admissible Mannheim mate of α . Thus, for all $s \in I$

$$\alpha^*(s) = \alpha(s) + \lambda(s)N(s). \quad (3.1)$$

The mate of an admissible Mannheim curve is denoted by (α, α^*) , see Figure 1.




Figure 1. The admissible Mannheim partner curves

Theorem 3.1 Let (α, α^*) be a mate of admissible Mannheim pair in G_3 . Then function λ is constant on I and defined by equation (3.1).

Proof. Let α be an admissible Mannheim curve in G_3 and α^* be an admissible Mannheim mate of α . Let the pair of $\alpha(s)$ and $\alpha^*(s)$ be of corresponding points of α and α^* . Then the curve $\alpha^*(s)$ is given by (3.1). Differentiating (3.1) with respect to s and using Frenet equations,

$$T^* \frac{ds^*}{ds} = T + \lambda' N + \lambda \tau B \quad (3.2)$$

is obtained.

Here and here after prime denotes the derivative with respect to s . Since N is coincident with B^* in the same direction, we have

$$\lambda'(s) = 0, \quad (3.3)$$

that is, λ is constant. This theorem proves that the distance between the curve α and its Mannheim mate α^* is constant at the corresponding points of them. It is notable that if α^* is an admissible Mannheim mate of α . Then α is also Mannheim mate of α^* because the relationship obtained in theorem between a curve and its Mannheim mate is reciprocal one. ■

Theorem 3.2 *Let α be an admissible curve with arc length parameter s . α is an admissible Mannheim curve if and only if the torsion τ of α is constant.*

Proof. Let (α, α^*) be a mate of an admissible Mannheim curves, then there exists the relation

$$\begin{aligned} T(s) &= \cos \theta T^*(s) + \sin \theta N^*(s) \\ B(s) &= -\sin \theta T^*(s) + \cos \theta N^*(s) \end{aligned} \quad (3.4)$$

and

$$\begin{aligned} T^*(s) &= \cos \theta T(s) - \sin \theta B(s) \\ N^*(s) &= \sin \theta T(s) + \cos \theta B(s) \end{aligned} \quad (3.5)$$

where θ is the angle between T and T^* at the corresponding points of $\alpha(s)$ and $\alpha^*(s)$, (see Figure 1).

By differentiating (3.5) with respect to s , we get

$$\tau^* B^* \frac{ds^*}{ds} = \frac{d(\sin \theta)}{ds} T + \sin \theta \kappa N + \cos \theta \tau N + \frac{d(\cos \theta)}{ds} B. \quad (3.6)$$

Since the principal normal vector field N of the curve α and the binormal vector field B^* of its Mannheim mate curve, then it can be seen that θ is a constant angle.

If we equations (3.2) and (3.5) is considered, then

$$\lambda \tau \cot \theta = 1 \quad (3.7)$$

is obtained. According to Theorem 3.1 and constant angle θ , $u = \lambda \cot \theta$ is constant. Then from equation (3.7), $\tau = \frac{1}{u}$ is constant, too.

Hence the proof is completed. ■

Theorem 3.3 (Schell's Theorem). *Let (α, α^*) be a mate of an admissible Mannheim curves with torsions τ and τ^* , respectively. The product of torsions τ and τ^* is constant at the corresponding points $\alpha(s)$ and $\alpha^*(s)$.*

Proof. Since α is an admissible Mannheim mate of α^* , then (3.1) also can be given by

$$\alpha = \alpha^* - \lambda B^*. \quad (3.8)$$

By taking differentiation of last equation and using equation (3.4),

$$\tau^* = \frac{1}{\lambda} \tan \theta \quad (3.9)$$

can be given. By the helps of (3.7), the equation below is obtained easily;

$$\tau \tau^* = \frac{\tan^2 \theta}{\lambda^2} = \text{constant} \quad (3.10)$$

This completes the proof. ■

Theorem 3.4 *Let (α, α^*) be an admissible Mannheim mate with curvatures κ, κ^* and torsions τ, τ^* of α and α^* , respectively. Then their curvatures and torsions satisfy the following relations*

$$\begin{aligned} i) \quad & \kappa^* = -\frac{d\theta}{ds^*} \\ ii) \quad & \kappa = \tau^* \frac{ds^*}{ds} \sin \theta \\ iii) \quad & \tau = -\tau^* \frac{ds^*}{ds} \cos \theta. \end{aligned}$$

Proof. *i)* Let us consideration equation (3.4), then we have

$$\langle T, T^* \rangle = \cos \theta. \quad (3.11)$$

By differentiating last equation with respect to s^* and using the Frenet equations of α and α^* , we reach

$$\langle \kappa(s)N(s)\frac{ds}{ds^*}, T^*(s) \rangle + \langle T(s), \kappa^*(s)N^*(s) \rangle = -\sin \theta \frac{d\theta}{ds^*}. \quad (3.12)$$

Since the principal normal N of α and binormal B^* of α^* are linearly dependent. By considering equations (3.4) and (3.12), we reach

$$\kappa^*(s) = -\frac{d\theta}{ds^*}. \quad (3.13)$$

If we take into consideration $\langle T, B^* \rangle$, $\langle B, B^* \rangle$ scalar products and (2.5), (3.4), (3.5) equations, then we can easily prove *ii)* and *iii)* items of the theorem, respectively.

The relations given in *ii)* and *iii)* of the last theorem, we obtain

$$\frac{\kappa}{\tau} = -\tan \theta = \text{constant}.$$

■

References

- [1] Liu, H. and Wang, F., *Mannheim partner curves in 3-space*, Journal of Geometry, 88 (2008), 120-126.
- [2] Orbay, K. and Kasap, E., *On Mannheim partner curves in E^3* , International Journal of Physical Sciences, 4 (5) (2009), 261-264.
- [3] Röschel, O., *Die Geometrie des Galileischen Raumes*, Habilitationsschrift, Leoben, (1984).
- [4] Sipus, Z. M., *Ruled Weingarten surfaces in the Galilean space*, Periodica Mathematica Hungarica, 56 (2) (2008), 213-225.
- [5] Kamenarovic, I., *Existence theorems for ruled surfaces in the Galilean space G_3* , Rad Hrvatskeakad. Znan. Umj. Mat. 10 (1991), 183-196.
- [6] Pavkovic, B. J. and Kamenarovic, I., *The equiform differential geometry of curves in the Galilean space G_3* , Glasnik Matematicki, 22 (42) (1987), 449-457.
- [7] Öğrenmiş, A. O., Bektaş, M. and Ergüt, M., *The characterizations for helices in the Galilean space G_3* , Int. J. of Pure and Appl. Math., 16 (1) (2004), 31-36.
- [8] Öğrenmiş, A. O., Ergüt, M. and Bektaş, M., *On the helices in the Galilean space G_3* , Iranian J. of Sci. Tec, Trans. A., 31 (2) (2007), 177-181.
- [9] Molnar, E., *The projective interpretation of the eight 3-dimensional homogeneous geometries*, Beitrage zur Algebra und Geometrie Contributions to Algebra and Geometry 38 (2) (1997), 261-288.